

AD-A126 908

THE MOLECULAR WEIGHT DISTRIBUTION FOR MOLECULES  
REQUIRING ROTATIONAL DIFF..(U) MASSACHUSETTS INST OF  
TECH CAMBRIDGE DEPT OF MATERIALS SCIENC.. L H PEEBLES

1/1


UNCLASSIFIED

OCT 82 TR-15 N00014-75-C-0542

F/G 20/8

NL

END  
DATE  
FILMED  
5 83  
DTIC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

AD A 126908

OFFICE OF NAVAL RESEARCH

Contract N00014-75-C-0542

Task No. NR 356-534

Technical Report No. 15 ✓

The Molecular Weight Distribution for Molecules Requiring  
Rotational Diffusion Prior to Reaction

by

L. H. PEEBLES, JR.

Prepared for Publication in MACROMOLECULES

Office of Naval Research  
Detachment, Boston  
495 Summer Street  
Boston, MA 02210

and

Department of Materials Science and Engineering  
Massachusetts Institute of Technology  
Cambridge, MA 02139

October 1982

Reproduction in whole or in part is permitted for  
any purpose of the United States Government

This document has been approved for public release  
and sale; its distribution is unlimited

\*Current address: Office of Naval Research  
800 N. Quincy Street  
Arlington, VA 22217

DTIC FILE COPY

83 04 15 115

DTIC  
ELECTE  
APR 18 1983  
B

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report #15	2. GOVT ACCESSION NO. AD A126908	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) The Molecular Weight Distribution for Molecules Requiring Rotational Diffusion Prior to Reaction		5. TYPE OF REPORT & PERIOD COVERED
7. AUTHOR(s) L. H. Peebles, Jr.		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Massachusetts Institute of Technology Cambridge, MA 02139		8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0542
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research, Code 413 800 N. Quincy Street Arlington, VA 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 356-534
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE October 1982
		13. NUMBER OF PAGES 18
		15. SECURITY CLASS. (of this report)
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited; <del>reproduction in whole or in part is permitted for any purpose of the United States government.</del>		
17. DISTRIBUTION STATEMENT (of the abstract entered in block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  Molecular weight distribution; diffusion; scaling law		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The molecular weight distributions have been calculated for rigid rods which must rotate into a collinear conformation prior to reaction. The underlying assumptions include bimolecular reaction without by-product formation and a rotary diffusion parameter that is proportional to molecular length through the term $(i^{-N} + j^{-N})$ where $i$ and $j$ are rod lengths and $N$ is an arbitrary parameter. To reduce the number of simultaneous differential equations that must be solved, an empirical scaling law is introduced which is independent of chain length. Essentially asymptotic values are obtained for number (over)		

DD FORM 1473  
1 JAN 73EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-LF-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

## Introduction

↓

In 1939 Flory announced the principle of equal reactivity of polymer chains (1). The principle maintains that the ability of a chain molecule to form larger molecules is independent of the chain length. The argument was very significant at that time for two reasons: First, the existence of high molecular weight polymers was considered by some to be improbable. These proponents argued that the high molecular weight species must have a higher diffusion constant, relative to the lower molecular weight species, which would eventually limit the extent of polymerization. Studies on condensation-type polymers with random coils demonstrated the general validity of the principle of equal reactivity (2). The second significant result was that the distribution of molecular weights could be readily calculated from kinetic parameters that were independent of chain length.

↗


The major significant deviations from this principle seem to occur only with very low molecular weight species where inductive or polarization effects can be transmitted across a small molecule. Indeed, Lenz (3) cites a number of cases where the reactivity of a functional group on a difunctional monomer depends on whether or not the other functional group has reacted. Challa (4) showed a similar effect on the rate of polymerization of poly-(ethylene terephthalate) where the reactivity of the "monomer" bis-( $\beta$ -hydroxyethyl) terephthalate differs from that of higher oligomers and polymer. Peebles and Wagner (5) examined the transesterification kinetics of dimethyl terephthalate and ethylene glycol in the presence of a catalyst.

They concluded that the second transesterification reaction was some three times faster than the first transesterification reaction. The overall transesterification of the terephthalate monomer must somehow involve some type of molecular transition state containing the catalyst because attempts to verify the faster transesterification reaction with the partly transesterified material resulted in absolutely no reaction (6). Thus, the documented "violations" of the principle of equal reactivity seems to be restricted to very small chain segments of random coil molecules.

Several calculations have been made to determine how departures from the principle of equal reactivity would influence the molecular weight distribution of a polymer. Nanda and Jain (7) assumed that the second order reaction constant for a condensation polymer was proportional to the molecular weight of the formed polymer through the relation  $k(1 + \frac{6}{5}i)$ , where  $k$  and  $\frac{6}{5}$  are constants and  $i$  is the number of monomers in a chain of length  $i$ ,


to obtain an analytical solution for the molecular weight distribution. Magat (8) developed simultaneous integral equations for stepwise polymerizations without termination for a number of conditions where the rate constant for propagation varied with chain length. His approaches are limited, however, in that the precise polymerization conditions must be specified before the equations can be evaluated.

A new series of stiff rod-like polymers (9) have recently been synthesized and characterized with a repeating unit of the form



where the "X" atom can be oxygen, sulfur, or singly hydrated nitrogen.

# The condensation of the functional groups



appears to occur simultaneously to form the benzobisoxazole because intermediate functionalities have not been detected by chemical or spectral means (9e). These observations suggest that the interacting molecules must be collinear (or nearly so) before reaction can take place. Furthermore, once reaction has commenced, the entire series of reactions to form the cyclic structure must occur quite rapidly. If such is the case, then the rate determining step is the ability of rods to rotate into a collinear configuration rather than the coupling step. Here the principle of equal reactivity is not violated; the reactivity of each end is independent of the length of the chain to which it is attached. The purpose of this paper is to calculate the molecular weight distribution for randomly oriented rod-like molecules which must diffuse or rotate into a collinear configuration prior to the coupling reaction.



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	

### Mathematical Development

The reaction between molecules of size  $i$  reacting with molecules of size  $j$  to form a molecule of size  $i + j$  irreversibly and without by-product



is postulated to have a rate of formation proportional to  $k c_{ij} x_i x_j$  where  $k$  is the specific second order rate constant for coupling,  $c_{ij}$  is a function of the molecular lengths  $i$  and  $j$  and is a measure of the ability of molecules to become collinear, and  $x_i, x_j$  are the concentrations of molecules of size  $i$  and  $j$ .

To simplify the notation, the constant  $k$  is adsorbed into the time derivative.

Thus from eq (1), the rate of formation of  $x_i$  is given by

$$\dot{x}_i = \sum_{j=1}^{i-1} c_{j,i-j} x_j x_{i-j} - x_i \sum_{j=1}^{\infty} c_{ij} x_j \quad (2)$$

The monomer  $x_1$  is considered to react at the rate

$$\dot{x}_1 = - x_1 \sum_{j=1}^{\infty} c_{1j} x_j \quad (3)$$

These rate formulas can be summed to give the following time derivative functions

$$\sum_{i=1}^{\infty} i \dot{x}_i = 0 \quad \text{line up?} \quad (4)$$

since no by-product is formed.

$$\sum_{i=1}^{\infty} i \dot{x}_i = - \sum_{i=1}^{\infty} x_i \sum_{j=1}^{\infty} c_{ij} x_j \quad \text{line up?} \quad (5)$$

$$\sum_{i=1}^{\infty} i^2 \dot{x}_i = \sum_{i=1}^{\infty} i x_i \sum_{j=1}^{\infty} j c_{ij} x_j \quad (6)$$



If the variable  $c_{ij}$  can be specified, then the molecular weight distribution and the moments of the distribution can in principal be determined by computer calculations.

Because many polymers have degrees of polymerization in excess of 100, and perhaps into the thousands or millions, it is not reasonable to solve the excessive number of simultaneous differential equations given by (2) and (3). However, if some type of scaling law can be found which will permit calculation of the distribution for a low number average molecular weight distributions, then computer calculations are feasible.

The generalized exponential distribution function

$$F(i, m, k) = m y^{k/m} i^{k-1} [\exp(-y i^m)] / P(k/m) \quad (7) \quad \text{G. k cap P}$$

$$W(i, m, k) = m y^{(k+1)/m} i^k [\exp(-y i^m)] / P[(k+1)/m] \quad (8)$$

where  $\langle i \rangle_n = \Gamma[(k+1)/m] / y^{1/m} P(k/m)$ ,

can be rewritten in a scaling law form with  $i/\langle i \rangle_n$  as the independent variable.

$$F(i, m, k) \langle i \rangle_n = \frac{m y^k}{P(k/m)} \left[ \frac{i}{\langle i \rangle_n} \right]^{k-1} \exp \left[ - (g i / \langle i \rangle_n)^m \right] \quad (9)$$

$$W(i, m, k) \langle i \rangle_n = \frac{m y^k}{P(k/m)} \left[ \frac{i}{\langle i \rangle_n} \right]^k \exp \left[ - (g i / \langle i \rangle_n)^m \right] \quad (10)$$

with  $g = \{P[(k+1)/m]\} / P(k/m)$

This distribution is a generalized form of the Schulz distribution ( $m = 1$ ),

the Tung-Weibull distribution ( $k = m-1$ ,  $k > 0$ ), and a good approximation of the most probable distribution ( $m = k = 1$ ) at large values of  $\langle i \rangle_n$ .

The next question to consider is the form of the function  $c_{ij}$  which determines the ability of the molecules to rotate into a collinear conformation. Doe and Edwards (9) suggest that the rotational diffusion constant of a rod

depends on  $(\ln i)/i^3$  in dilute solution and higher inverse powers of  $i$  in more concentrated solutions, whereas Bird (10) suggests that the rotation of a rigid dumbbell depends on the square of the length ( $i^2$ ). The assumption can then be made that if rod  $j$  were held fixed in space while rod  $i$  was allowed to rotate freely, then the rate of rotation would be proportional to  $j^{-N}$ , where  $N$  is an adjustable parameter. If rod  $j$  were held fixed and rod  $i$  allowed to rotate, the rate is proportional to  $i^{-N}$ . The overall average rate is then

$$c_{ij} = (i^{-N} + j^{-N})/2 \quad (11)$$

This equation can then be substituted into the set of simultaneous differential equations 2 <sup>-6</sup>. The initial concentration of monomer,  $x_1$  was selected to be unity. Initially,  $x_1$  was taken as the independent variable. A set of  $K$  simultaneous differential equations  $\dot{x}_i/\dot{x}_{ind}$ , where  $\dot{x}_{ind}$  is the change in concentration with time of the independent variable, was set up for solution by a computer. The value of  $K$  was selected to be large enough so that  $\sum i x_i > .999$  over the range of integration considered. A Runge-Kutta method was used to obtain values of  $x_i$ , from  $i = ind$  to  $i = K$ . After each integration, an interger value of  $L = 0.8K$  was tested to see whether  $Lx_L < 0.0005$ . If the test failed, a larger value of  $K$  was selected, the program retreated to the results obtained on a previously performed integration, and computation continued. This procedure allowed only a small number of equations to be solved at the beginning of the solution, but as  $\langle i \rangle_n$  increased, more equations were solved, and hence more computer time required. Also, as  $\langle i \rangle_n$  increased, values of  $x_{ind}$  decreased and eventually became too small to permit reliable values of  $\dot{x}_i/\dot{x}_{ind}$  to be computed as evidenced by fluctuations in  $x_{ind}$ . As  $x_{ind}$  approached this limiting value, a new independent variable was selected and concentrations of all  $i$  variables below  $x_{ind-1}$  were arbitrarily set to zero. Extensive testing of this procedure by choosing various values of  $i$  for the independent variable exhibited no effect on the calculated distribution provided that  $x_{ind} < x_{max}$ .

### Results

Integration of eq's 2, 3, 5 and 6 were performed for integer values of  $N$  in eq 11 as a function of conversion. Values of  $\langle i \rangle_n$  varied from unity (initial condition) to  $\langle i \rangle_n$  values in excess of 50. From the  $x_i$  values so computed, values of  $x_i \langle i \rangle_n / \sum x_i$  were obtained. For all values of  $N$  used, except for  $N = 0$  (the most probable distribution) the scaling law held to within the precision of the data when  $\langle i \rangle_n$  values exceeded ten. The results are presented in Fig.'s 1 and 2 for the distribution as a function of  $i/\langle i \rangle_n$  and in Fig. 3 for the heterogeneity index  $\langle i \rangle_w / \langle i \rangle_n$  as a function of conversion.

### Discussion

The empirical observation that the scaling laws  $F(i) \langle i \rangle_n$  and  $W(i) \langle i \rangle_n$  are functions only of  $i/\langle i \rangle_n$  means that inexpensive computer calculations can be performed to evaluate these functions at small values of  $\langle i \rangle_n$ .  
then  
The function can be used directly to evaluate the molecular weight distributions for large values of  $\langle i \rangle_n$ . This technique can be used for other schemes relating kinetics to reactor designs, etc. where numerical calculations are required.

When the value of  $N$  in equation 11 is nonzero, the rate of polymerization  $\Sigma x_i$ , eq 5, decreases with increasing conversion. As  $N$  increases, the rate of polymerization decreases at a faster rate. Both the scaled mole fraction and weight fraction distributions show a maximum in the range  $i/\langle i \rangle_n$  of 0.6-0.7, the maximum shifting to higher values with increasing  $N$ . The breadth of the distribution, however, becomes narrower, and reaches an asymptotic value at  $\langle i \rangle_n$  values of approximately 10. The application of these calculations to the observed distributions of Poly([benzo(1,2-d:5,4-d') bisoxazole-2,6 -diyl]-1,4-phenylene) has been discussed by Cotts and Berry (9e).

### Acknowledgment

The author wishes to express his gratitude to Dr. Guy Berry who suggested the problem, to the Office of Naval Research for partial support of the work, and to the Department of Materials Science and Engineering, Massachusetts Institute of Technology, for an appointment as Visiting Scientist.

### References and Notes

1. P. J. Flory, J. Am. Chem. Soc. 1936, 58, 1877
2. P. J. Flory, J. Am. Chem. Soc. 1939, 61, 3334; 1940, 62, 347
3. R. W. Lenz, "Organic Chemistry of Synthetic High Polymers" Interscience, New York, N.Y., 1967, p.
4. G. Challa, Makromol. Chem., 1960, 38, 105
5. L. H. Peebles, Jr. & W. S. Wagner, J. Phys. Chem, 1959, 63, 1206
6. Recent investigations by Wagner et al, private communication, 1981, suggest that the catalyst-monomer system follows third order kinetics, which tends to support involvement of a complex molecular transition state (5).
7. V. S. Nanda and S. C. Jain, J. Chem. Phys., 1968, 49, 1318
8. H. J. R. Maget, J. Polym. Sci, 1964, A2, 1281
9. a. J. F. Wolfe & F. E. Arnold, Macromolecules, 1981, 14, 909;  
b. F. J. Wolfe, B. H. Loo, F. E. Arnold ibid, p 915;  
c. E. W. Choe & S. N. Kim, ibid, p 920;  
d. R. E. Evers, F. E. Arnold, and T. E. Helminiak, ibid, p 925;  
e. D. B. Cotts & G. C. Berry, ibid, p 930;  
f. M. W. Wellman, W. W. Adams, R. A. Wolfe, D. S. Dudis, D. R. Wiff, & A. V. Fratini, ibid, p 935;  
g. S. -G. Chu, S. Venkatraman, G. C. Berry and Y. Einaga, ibid, p 939;  
h. W. J. Welsh, D. Bhavmik, and J. E. Mark, ibid, 947;  
i. D. B. Laumik, W. J. Welsh, H. H. Jaffe, & J. E. Mark, ibid, 951;  
j. R. R. Matheson, Jr. & P. J. Flory, ibid, 954;  
k. L. D. Bhavmik, H. H. Jaffe & J. E. Mark, ibid, 1125;  
l. S. R. Allen, A. G. Filippov, R. J. Farris, E. L. Thomas, C. P. Wong, G. C. Berry, & E. C. Chenevey, ibid, 1135.
10. M. Doi & S. F. Edwards, J. Chem. Soc., Faraday Trans II, 1978, 74, 918
11. R. B. Bird, O. Hassager, R. C. Armstrong, and C. F. Curtis, Dynamics of Polymer Liquids, Vol II, Chapt 11. John Wiley & Sons, New York 1977.

Legend for Figures

Figure 1. The scaled mole fraction of polymer as a function of  $i/\langle i \rangle_n$  for various values of  $N$  in eq 11

Figure 2. The scaled weight fraction of polymer as a function of  $i/\langle i \rangle_n$  for various values of  $N$  in eq 11

Figure 3. The heterogeneity index  $\langle i \rangle_n / \langle i \rangle_n$  as a function of  $\langle i \rangle_n$  and  $N$ .




Fig 1




Fig 2






Fig 3

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code -13 800 North Quincy Street Arlington, Virginia 22217	2	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
ONR Pasadena Detachment Attn: Dr. R. J. Marcus 1000 East Green Street Pasadena, California 91106	1	Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Dean William Tolles Naval Postgraduate School Monterey, California 93940	1
Dr. Fred Saalfeld Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
U.S. Army Research Office Attn: CRD-AA-IP P. O. Box 12211 Research Triangle Park, N.C. 27709	1	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Mr. Vincent Schaper DTNSRDC Code 1803 Annapolis, Maryland 21402	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19102	1
Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1	Mr. A. M. Ansalone Administrative Librarian PLASTECARRADCOM Bldg 3401 Dover, New Jersey 07801	1

Copy available to DTIC does not  
permit fully legible reproduction

TECHNICAL REPORT DISTRIBUTION LIST, 356A

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. M. Broadhurst Bulk Properties Section National Bureau of Standards U. S. Department of Commerce Washington, D.C. 20234	2	Dr. K. D. Pae Department of Mechanics and Materials Science Rutgers University New Brunswick, New Jersey 08903	1
Naval Surface Weapons Center Attn: Dr. J. M. Augl, Dr. B. Hartman White Oak Silver Spring, Maryland 20910	1	NASA-Lewis Research Center Attn: Dr. T. T. Serofini, MS-49-1 2100 Brookpark Road Cleveland, Ohio 44135	1
Dr. G. Goodman Globe Union Incorporated 5757 North Green Bay Avenue Milwaukee, Wisconsin 53201	1	Dr. Charles H. Sherman Code TD 121 Naval Underwater Systems Center New London, Connecticut 06320	1
Professor Hatsuo Ishida Department of Macromolecular Science Case-Western Reserve University Cleveland, Ohio 44106	1	Dr. William Risen Department of Chemistry Brown University Providence, Rhode Island 02191	1
Dr. David Soong Department of Chemical Engineering University of California Berkeley, California 94720	1	Mr. Robert W. Jones Advanced Projects Manager Hughes Aircraft Company Mail Station D 132 Culver City, California 90230	1
Dr. Curtis W. Frank Department of Chemical Engineering Stanford University Stanford, California 94035	1	Dr. C. Giori IIT Research Institute 10 West 35 Street Chicago, Illinois 60616	1
Picatinny Arsenal Attn: A. M. Anzalone, Building 3401 SMUPA-FR-M-D Dover, New Jersey 07801	1	Dr. R. S. Roe Department of Materials Science and Metallurgical Engineering University of Cincinnati Cincinnati, Ohio 45221	1
Dr. J. K. Gillham Department of Chemistry Princeton University Princeton, New Jersey 08540	1	Dr. Robert E. Cohen Chemical Engineering Department Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1
Dr. E. Baer Department of Macromolecular Science Case Western Reserve University Cleveland, Ohio 44106	1	Dr. T. P. Conlon, Jr., Code 3622 Sandia Laboratories Sandia Corporation Albuquerque, New Mexico	1

TECHNICAL REPORT DISTRIBUTION LIST, 356A

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. Martin Kaufman Code 38506 Naval Weapons Center China Lake, California 93555	1	Professor C. S. Paik Sung Department of Materials Sciences and Engineering Room 8-109 Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1
Professor S. Senturia Department of Electrical Engineering Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1	Professor Brian Newman Department of Mechanics and Materials Science Rutgers, The State University Piscataway, New Jersey 08854	1
Dr. T. J. Reinhart, Jr., Chief Composite and Fibrous Materials Branch Nonmetallic Materials Division Department of the Air Force Air Force Materials Laboratory (AFSC) Wright-Patterson AFB, Ohio 45433	1	Dr. John Lundberg School of Textile Engineering Georgia Institute of Technology Atlanta, Georgia 30332	1
Dr. J. Lando Department of Macromolecular Science Case Western Reserve University Cleveland, Ohio 44106	1	Dr. D. R. Uhlmann Department of Materials Science Massachusetts Institute of Technology Cambridge, MA 02139	1
Dr. J. White Chemical and Metallurgical Engineering University of Tennessee Knoxville, Tennessee 37916	1		
Dr. J. A. Manson Materials Research Center Lehigh University Bethlehem, Pennsylvania 18015	1		
Dr. R. F. Helmreich Contract RD&E Dow Chemical Co. Midland, Michigan 48640	1		
Dr. R. S. Porter Department of Polymer Science and Engineering University of Massachusetts Amherst, Massachusetts 01002	1		
Professor Garth Wilkes Department of Chemical Engineering Virginia Polytechnic Institute and State University Blacksburg, Virginia 24061	1		

